

On some p -adic power series attached to the arithmetic of $\mathbb{Q}(\zeta_p)$

Bruno Anglès
 Université de Caen,
 Laboratoire Nicolas Oresme, CNRS UMR 6139,
 Campus II, Boulevard Maréchal Juin,
 BP 5186, 14032 Caen Cedex, France.
 E-mail: angles@math.unicaen.fr

October 11 2005

Let p be a prime number, $p \geq 5$. Let θ be an even and non-trivial character of $\text{Gal}(\mathbb{Q}(\mu_p)/\mathbb{Q})$. Let $f(T, \theta) \in \mathbb{Z}_p[[T]]$ be the Iwasawa power series associated to the p -adic L-function $L_p(s, \theta)$ (see [12]), i.e.:

$$\forall n \geq 1, n \equiv 0 \pmod{p-1}, f((1+p)^{1-n} - 1, \theta) = L(1-n, \theta),$$

where $L(s, \theta)$ is the usual Dirichlet L-series. In 1979, in their celebrated article [4], Ferrero and Washington have proved that (see also [1]):

$$f(T, \theta) \not\equiv 0 \pmod{p}.$$

Thus we can write:

$$f(T, \theta) \equiv T^{\lambda(\theta)} \bar{U}(T) \pmod{p},$$

where $\bar{U}(T) \in \mathbb{F}_p[[T]]^*$. The Iwasawa lambda-invariant $\lambda(\theta)$ is not well-understood. By a heuristic argument due to Ferrero and Washington, one could expect that for sufficiently large p (see [5]):

$$\sum_{\theta \text{ even}, \theta \neq 1} \lambda(\theta) \leq \frac{\text{Log}(p)}{\text{Log}(\text{Log}(p))}.$$

Furthermore, if $p < 4000000$, we have $\lambda(\theta) \leq 1$, and one would reasonably expect (see [8]):

$$\lambda(\theta) < p.$$

The only known result is due to Ferrero and Washington ([4]): for sufficiently large p we have:

$$\lambda(\theta) \leq p^{\text{Log}(p)^{4\varphi(p-1)^4}}.$$

Now observe that:

$$f'(T, \theta) \equiv \lambda(\theta)T^{\lambda(\theta)-1}\bar{U}(T) + T^{\lambda(\theta)}\bar{U}'(T) \pmod{p}.$$

Thus if $\lambda(\theta) \geq 1$ and $\lambda(\theta) \not\equiv 0 \pmod{p}$, we have that $f'(T, \theta) \not\equiv 0 \pmod{p}$.

The aim of this paper is to prove that for all θ even, $\theta \neq 1$, we have (Corollary 4.4):

$$f'(T, \theta) \not\equiv 0 \pmod{p}.$$

This fact comes from properties of some power series that are connected to polynomials introduced by Mirimanoff at the beginning of the XXth century.

1 Notations

Let p be a prime number, $p \geq 5$. Let $\overline{\mathbb{Q}_p}$ be an algebraic closure of $\mathbb{Q}_p$. All the extensions of $\mathbb{Q}_p$ considered in this paper are contained in $\overline{\mathbb{Q}_p}$. Let v_p be the p -adic valuation on $\overline{\mathbb{Q}_p}$ such that $v_p(p) = 1$. We denote the Iwasawa p -adic logarithm on $\overline{\mathbb{Q}_p}$ by Log_p .

If A is a commutative ring, we denote the set of invertible elements of A by A^* .

For every integer d , $d \geq 1$, set $\mu_d = \{z \in \overline{\mathbb{Q}_p} \mid z^d = 1\}$. If $\rho \in \cup_{d \geq 1} \mu_d$, we denote the order of ρ by $o(\rho)$. Set $\mu_{p^\infty} = \cup_{n \geq 0} \mu_{p^{n+1}}$. For all $n \geq 0$, let $\zeta_{p^{n+1}} \in \mu_{p^{n+1}}$ such that: $\zeta_p \neq 1$ and $\forall n \geq 0$, $\zeta_{p^{n+2}}^p = \zeta_{p^{n+1}}$.

Let:

- $K_n = \mathbb{Q}_p(\mu_{p^{n+1}})$,
- $O_n = \mathbb{Z}_p[\zeta_{p^{n+1}}]$,

- $\pi_n = \zeta_{p^{n+1}} - 1$,
- $U_n = 1 + \pi_n O_n$,
- $\Gamma_n = \text{Gal}(K_n/K_0)$,
- $\Delta = \text{Gal}(K_0/\mathbb{Q}_p)$,
- $K_\infty = \mathbb{Q}_p(\mu_{p^\infty})$,
- $\Gamma = \text{Gal}(K_\infty/K_0)$.

Let $\gamma_0 \in \Gamma$ be such that $\forall \zeta \in \mu_{p^\infty}, \zeta^{\gamma_0} = \zeta^{1+p}$.

For $a \in \mathbb{Z}_p^*$, we write:

$$a = \omega(a) < a >,$$

where ω is the Teichmüller character (i.e. $\omega(a) = \lim_n a^{p^n} \in \mu_{p-1}$) and $< a > \equiv 1 \pmod{p}$.

Let $n \geq 0$. For $a \in \mathbb{Z}_p^*$, let $\sigma_n(a) \in \text{Gal}(K_n/\mathbb{Q}_p)$ be such that:

$$\zeta_{p^{n+1}}^{\sigma_n(a)} = \zeta_{p^{n+1}}^a.$$

We have $\sigma_n(a) = \delta(a)\gamma_n(a)$, where $\delta(a) \in \Delta$ and $\gamma_n(a) \in \Gamma_n$. Observe that:

$$\zeta_{p^{n+1}}^{\gamma_n(a)} = \zeta_{p^{n+1}}^{<a>}.$$

For $a \in \mathbb{Z}_p^*$, let $\sigma(a) \in \Gamma \times \Delta$ be such that:

$$\forall \zeta \in \mu_{p^\infty}, \zeta^{\sigma(a)} = \zeta^a.$$

Note that $\Delta = \{\sigma(\omega(a)) \mid a = 1, \dots, p-1\}$. For $\theta \in \widehat{\Delta} = \text{Hom}(\Delta, \mu_{p-1})$, we set:

$$e_\theta = \frac{1}{p-1} \sum_{\delta \in \Delta} \theta^{-1}(\delta) \delta \in \mathbb{Z}_p[\Delta].$$

Let T be an indeterminate over $\overline{\mathbb{Q}_p}$. Let L be a finite extension of $\mathbb{Q}_p$. Let O_L be the valuation ring of L . The restriction maps $\text{Res}_{n+1,n} : \Gamma_{n+1} \rightarrow \Gamma_n$ induce maps $\text{Res}_{n+1,n} : O_L[\Gamma_{n+1}] \rightarrow O_L[\Gamma_n]$. Thus we can take the inverse limit of the group rings $O_L[\Gamma_n]$ with respect to these maps, and by [12], Theorem 7.1, we have:

$$O_L[[T]] \simeq \varprojlim O_L[\Gamma_n],$$

where T corresponds to $\gamma_0 - 1$. We set $\Lambda_L = O_L[[T]]$ and $\Lambda = \Lambda_{\mathbb{Q}_p}$. For all $n \geq 0$, we set $\omega_n(T) = (1+T)^{p^n} - 1$. Recall that:

$$\forall n \geq 0, \frac{\Lambda_L}{\omega_n(T)\Lambda_L} \simeq O_L[\Gamma_n].$$

Note that for all $n \geq 0$, $N_{K_{n+1}/K_n}(U_{n+1}) \subset U_n$, where N_{K_{n+1}/K_n} is the norm map from K_{n+1} to K_n . We denote the inverse limit of the principal units U_n with respect to the norm maps by U_∞ . Note that U_∞ is a $\Lambda[\Delta]$ -module.

2 Logarithmic derivatives

Let $u = (u_n)_{n \geq 0}$ be an element in U_∞ , recall that:

$$\forall n \geq 0, N_{K_{n+1}/K_n}(u_{n+1}) = u_n.$$

There exists a unique element $f_u(T) \in \Lambda$ such that ([12], Theorem 13.38, see also [3]):

$$\forall n \geq 0, f_u(\pi_n) = u_n.$$

Furthermore, if $x = \sum b_a \sigma(a) \in \mathbb{Z}_p[\Gamma \times \Delta]$ and if $u \in U_\infty$, then we have ([12], Lemma 13.48):

$$f_{u^x}(T) = \prod f_u((1+T)^a - 1)^{b_a}.$$

For $u \in U_\infty$ and for $n \geq 0$, we set:

$$D_n(u) = \zeta_{p^{n+1}} \frac{f'_u(\pi_n)}{f_u(\pi_n)} \in O_n.$$

We call the map $D_n : U_\infty \rightarrow O_n$ the n th logarithmic derivative of Coleman.

Lemma 2.1

- i) Let $u \in U_\infty$, $\forall n \geq 0$, $D_n(u^{\omega_n(T)}) \equiv 0 \pmod{p^{n+1}}$.
- ii) $\forall \theta \in \hat{\Delta}$, $\forall u \in U_\infty$, $\forall n \geq 0$, $D_n(u^{e_\theta}) = e_{\theta\omega^{-1}} D_n(u)$.

Proof

i) We have:

$$f_{u^{\omega_n(T)}}(T) = \frac{f_u((1+T)^{(1+p)^{p^n}} - 1)}{f_u(T)}.$$

Thus:

$$(1+T) \frac{f'_{u^{\omega_n(T)}}(T)}{f_{u^{\omega_n(T)}}(T)} = (1+p)^{p^n} (1+T)^{(1+p)^{p^n}} \frac{f'_u((1+T)^{(1+p)^{p^n}} - 1)}{f_u((1+T)^{(1+p)^{p^n}} - 1)} - (1+T) \frac{f'_u(T)}{f_u(T)}.$$

Note that:

$$(1 + \pi_n)^{(1+p)^{p^n}} - 1 = \pi_n.$$

Therefore:

$$D_n(u^{\omega_n(T)}) = \zeta_{p^{n+1}}((1+p)^{p^n} - 1) \frac{f'_u(\pi_n)}{f_u(\pi_n)} \equiv 0 \pmod{p^{n+1}}.$$

ii) Let $a \in \mathbb{Z}_p^*$ and let $u \in U_\infty$. Observe that:

$$D_n(u^{\sigma(a)}) = a\sigma_n(a)(D_n(u)).$$

Let $u \in U_\infty$, we get:

$$D_n(u^{e_\theta}) = \frac{1}{p-1} \sum_{a=1}^{p-1} \theta^{-1}(a) D_n(u^{\sigma(\omega(a))}).$$

We have:

$$D_n(u^{e_\theta}) = \frac{1}{p-1} \sum_{a=1}^{p-1} \theta^{-1}(a) \omega(a) \sigma_n(\omega(a))(D_n(u)).$$

The Lemma follows. $\diamond$

Proposition 2.2 *Let $\theta \in \widehat{\Delta}$, $\theta \neq 1, \omega$. The map D_n gives rise to a morphism of $\mathbb{Z}_p[\Gamma_n]$ -modules:*

$$\phi_n^\theta : \frac{U_n^{e_\theta}}{U_n^{p^{n+1}e_\theta}} \rightarrow \frac{e_{\theta\omega^{-1}}O_n}{p^{n+1}e_{\theta\omega^{-1}}O_n},$$

where T acts on the right via $(1+p)(1+T) - 1$.

Proof We have ([12], Theorem 13.54):

$$U_n^{e_\theta} \simeq \frac{U_\infty^{e_\theta}}{U_\infty^{\omega_n(T)e_\theta}}.$$

Let $v \in U_n^{e_\theta}$, then there exists $u = (u_n)_{n \geq 0} \in U_\infty^{e_\theta}$ such that $v = u_n$. We set:

$$\phi_n^\theta(v) \equiv D_n(u) \pmod{p^{n+1}}.$$

By Lemma 2.1, we get a morphism of $\mathbb{Z}_p$ -modules:

$$\phi_n^\theta : \frac{U_n^{e_\theta}}{U_n^{p^{n+1}e_\theta}} \rightarrow \frac{e_{\theta\omega^{-1}}O_n}{p^{n+1}e_{\theta\omega^{-1}}O_n}.$$

Let $v \in U_n^{e_\theta}$, and let $u = (u_n)_{n \geq 0} \in U_\infty^{e_\theta}$ such that $v = u_n$. We have:

$$\phi_n^\theta(u^T) \equiv D_n(u^{\gamma_0^{-1}}) \pmod{p^{n+1}}.$$

But:

$$D_n(u^{\gamma_0^{-1}}) \equiv ((1+p)\gamma_0 - 1)(D_n(u)) \pmod{p^{n+1}}.$$

The Proposition follows. $\diamond$

Theorem 2.3 *For all $n \geq 0$, set $T_n = \sum_{d=0}^n \zeta_{p^{d+1}}$. Let $\theta \in \widehat{\Delta}$, $\theta \neq 1, \omega$. Then:*

$$\forall n \geq 0, \text{Im}(\phi_n^\theta) = \frac{\mathbb{Z}_p[\Gamma_n]e_{\theta\omega^{-1}}T_n}{p^{n+1}e_{\theta\omega^{-1}}O_n}.$$

Proof Let Λ_n be the maximal order of $\mathbb{Q}_p[\Gamma_n]$. Then ([6], see also [7]):

$$O_n = \Lambda_n[\Delta] T_n.$$

Now observe that $p^{n+1}\Lambda_n \subset p\mathbb{Z}_p[\Gamma_n]$. Thus:

$$p^{n+1}e_{\theta\omega^{-1}}O_n \subset p\mathbb{Z}_p[\Gamma_n]e_{\theta\omega^{-1}}T_n.$$

Recall that there exists $\alpha \in \mu_{p-1} \setminus \{1\}$ such that ([12], Corollary 13.37):

$$\forall n \geq 0, U_n^{e_\theta} = \left(\frac{\alpha - \zeta_{p^{n+1}}}{\omega(\alpha - 1)} \right)^{\mathbb{Z}_p[\Gamma_n]e_\theta}.$$

For $\alpha \in \mu_{p-1} \setminus \{1\}$, set:

$$\rho_\infty(\alpha) = \left(\frac{\alpha - \zeta_{p^{n+1}}}{\omega(\alpha - 1)} \right)_{n \geq 0} \in U_\infty.$$

We have:

$$D_n(\rho_\infty(\alpha)) = \frac{-\zeta_{p^{n+1}}}{\alpha - \zeta_{p^{n+1}}}.$$

Thus $\text{Im}(\phi_n^\theta)$ is the $\mathbb{Z}_p[\Gamma_n]$ -module generated by the $e_{\theta\omega^{-1}} \frac{-\zeta_{p^{n+1}}}{\alpha - \zeta_{p^{n+1}}}$, $\alpha \in \mu_{p-1} \setminus \{1\}$. For $\alpha \in \mu_{p-1} \setminus \{1\}$, for all $n \geq 0$, set:

$$u_n(\alpha) = \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} \alpha^k \sigma_n(k) \in \mathbb{Z}_p[\Gamma_n \times \Delta].$$

We have:

$$\text{Res}_{n+1,n}(u_{n+1}(\alpha)) = \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+2}-1} \alpha^k \sigma_n(k).$$

Thus :

$$\text{Res}_{n+1,n}(u_{n+1}(\alpha)) = \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} \sigma_n(k) \left(\sum_{\ell=0}^{p-1} \alpha^{k+p^{n+1}\ell} \right).$$

Finally, we get:

$$\text{Res}_{n+1,n}(u_{n+1}(\alpha)) = u_n(\alpha).$$

Now, we have:

$$\frac{\alpha - 1}{\alpha \zeta_{p^{n+1}} - 1} - 1 = \frac{(\alpha \zeta_{p^{n+1}})^{p^{n+1}} - 1}{\alpha \zeta_{p^{n+1}} - 1} - 1.$$

Thus:

$$\frac{\alpha - 1}{\alpha \zeta_{p^{n+1}} - 1} - 1 = \sum_{k=1}^{p^{n+1}-1} \alpha^k \zeta_{p^{n+1}}^k.$$

Therefore:

$$\frac{\alpha - 1}{\alpha \zeta_{p^{n+1}} - 1} - 1 = \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} \alpha^k \sigma_n(k) (\zeta_{p^{n+1}}) + \sum_{k=1}^{p^n-1} \alpha^k \zeta_{p^n}^k.$$

Finally, we get:

$$\frac{\alpha - 1}{\alpha \zeta_{p^{n+1}} - 1} - 1 = u_n(\alpha)(T_n).$$

Now set:

$$u_n(\theta, \alpha) = \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} \theta(k) \omega^{-1}(k) \alpha^k \gamma_n(k).$$

We have:

$$(u_n(\theta, \alpha))_{n \geq 0} \in \Lambda.$$

Furthermore, we have:

$$e_{\theta\omega^{-1}} \frac{-\zeta_{p^{n+1}}}{\alpha - \zeta_{p^{n+1}}} = e_{\theta\omega^{-1}} \frac{1}{\alpha^{-1}\zeta_{p^{n+1}} - 1}.$$

Thus:

$$e_{\theta\omega^{-1}} \frac{-\zeta_{p^{n+1}}}{\alpha - \zeta_{p^{n+1}}} = \frac{1}{\alpha^{-1} - 1} u_n(\theta, \alpha^{-1})(e_{\theta\omega^{-1}} T_n).$$

Now:

$$\sum_{\alpha \in \mu_{p-1} \setminus \{1\}} u_0(\theta, \alpha) = (p-1)\theta(-1)\omega^{-1}(-1) \in \mathbb{Z}_p^*.$$

Thus there exists $\alpha \in \mu_{p-1} \setminus \{1\}$ such that for all $n \geq 0$, $u_n(\theta, \alpha) \in \mathbb{Z}_p[\Gamma_n]^*$.
The theorem follows. $\diamond$

We will also need the following Lemma:

Lemma 2.4 *Let Λ_n be the maximal order of $\mathbb{Q}_p[\Gamma_n]$. Under the isomorphism $\Lambda/\omega_n(T)\Lambda \simeq \mathbb{Z}_p[\Gamma_n]$, $p^{n+1}\Lambda_n$ corresponds to an ideal $\mathcal{U}_n$ of Λ such that $\omega_n(T)\Lambda \subset \mathcal{U}_n \subset (p, \omega_n(T))$ and $\lim_n \mathcal{U}_n = \{0\}$.*

Proof Set:

$$e_0 = \frac{1}{p^n} \sum_{\gamma \in \Gamma_n} \gamma,$$

and for $1 \leq d \leq n$, set:

$$e_d = \sum_{\chi \in \widehat{\Gamma_n}, f_\chi = p^{d+1}} e_\chi,$$

where the sum is over all the characaters of Γ_n of conductors p^{d+1} and :

$$e_\chi = \frac{1}{p^n} \sum_{\gamma \in \Gamma_n} \chi^{-1}(\gamma)\gamma.$$

We have:

$$p^{n+1}\Lambda_n = \oplus_{d=0}^n \mathbb{Z}_p[\Gamma_n] p^{n+1} e_d \subset p\mathbb{Z}_p[\Gamma_n].$$

Thus it is clear that $\omega_n(T)\Lambda \subset \mathcal{U}_n \subset (p, \omega_n(T))$. Now:

$$p^{n+1}e_0 \equiv p \left(\sum_{\ell=0}^{p^n-1} (1+T)^\ell \right) \pmod{\omega_n(T)}.$$

Thus:

$$p^{n+1}e_0 \equiv p \frac{\omega_n(T)}{T} \pmod{\omega_n(T)}.$$

Let d be an integer, $1 \leq d \leq n$. We have:

$$p^{n+1}e_d \equiv p \left(\sum_{\ell=0}^{p^n-1} (1+T)^\ell \left(\sum_{f_\chi=p^{d+1}} \chi(1+p)^\ell \right) \right) \pmod{\omega_n(T)}.$$

Thus:

$$p^{n+1}e_d \equiv p \left(\sum_{\ell=0}^{p^n-1} \text{Tr}_{K_{d-1}/\mathbb{Q}_p}(\zeta_{p^d}^\ell) (1+T)^\ell \right) \pmod{\omega_n(T)}.$$

Now recall that $\text{Tr}_{K_{d-1}/\mathbb{Q}_p}(\zeta_{p^d}^\ell) = 0$ if $v_p(\ell) < d-1$. Therefore:

$$p^{n+1}e_d \equiv p^d \left(\sum_{\ell=0}^{p^{n-d+1}-1} (1+T)^{\ell p^{d-1}} \text{Tr}_{K_0/\mathbb{Q}_p}(\zeta_p^\ell) \right) \pmod{\omega_n(T)}.$$

But:

$$\sum_{\ell=0, \ell \equiv 0 \pmod{p}}^{p^{n-d+1}-1} (1+T)^{\ell p^{d-1}} \text{Tr}_{K_0/\mathbb{Q}_p}(\zeta_p^\ell) = (p-1) \frac{\omega_n(T)}{\omega_d(T)},$$

and

$$\sum_{\ell=0, \ell \not\equiv 0 \pmod{p}}^{p^{n-d+1}-1} (1+T)^{\ell p^{d-1}} \text{Tr}_{K_0/\mathbb{Q}_p}(\zeta_p^\ell) = \frac{\omega_n(T)}{\omega_d(T)} - \frac{\omega_n(T)}{\omega_{d-1}(T)}.$$

Thus:

$$p^{n+1}e_d \equiv p^{d+1} \frac{\omega_n(T)}{\omega_d(T)} - p^d \frac{\omega_n(T)}{\omega_{d-1}(T)} \pmod{\omega_n(T)}.$$

The Lemma follows. $\diamond$

3 Mirimanoff's power series

Recall that Mirimanoff has introduced the following polynomials in $\mathbb{F}_p[T]$:

$$\forall j, 1 \leq j \leq p-1, \varphi_j(T) = \sum_{a=1}^{p-1} a^{j-1} T^a.$$

These polynomials have many beautiful properties and we refer the interested reader to [9], [11] and [2]. In this section, we will introduce some power series that are related to these polynomials.

Let L be a finite extension of $\mathbb{Q}_p$. Let π_L be a prime of L . Let $\theta \in \widehat{\Delta}$. Let $a \in O_L$ such that $a(a-1) \not\equiv 0 \pmod{\pi_L}$. For all $n \geq 0$, we set:

$$M_n(\theta, a) = \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} \frac{a^k}{a^{p^{n+1}}-1} \theta(k) \omega^{-1}(k) \gamma_n(k) \in O_L[\Gamma_n].$$

Lemma 3.1

$$(M_n(\theta, a))_{n \geq 0} \in \varprojlim O_L[\Gamma_n].$$

Proof We must prove that:

$$\forall n \geq 0, \text{Res}_{n+1,n}(M_{n+1}(\theta, a)) = M_n(\theta, a).$$

Now observe that:

$$\text{Res}_{n+1,n}(M_{n+1}(\theta, a)) = \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} \theta(k) \omega^{-1}(k) \gamma_n(k) \frac{1}{a^{p^{n+2}}-1} \left(\sum_{\ell=0}^{p-1} a^{k+\ell p^{n+1}} \right).$$

The Lemma follows. $\diamond$

By the above Lemma, $(M_n(\theta, a))_{n \geq 0}$ corresponds to a power series $M(T, \theta, a) \in \Lambda_L$. If $\theta = \omega^j$, $1 \leq j \leq p-1$, observe that:

$$M(0, \theta, a) \equiv \frac{\varphi_j(a)}{a^p-1} \pmod{\pi_L}.$$

Therefore we call $M(T, \theta, a)$ the Mirimanoff's power series attached to θ and a .

Lemma 3.2

$$M(T, \theta, a) = -\theta(-1) \omega^{-1}(-1) M(T, \theta, a^{-1}).$$

Proof

We have:

$$M_n(\theta, a) = \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} \frac{a^{p^{n+1}-k}}{a^{p^{n+1}}-1} \theta(p^{n+1}-k) \omega^{-1}(p^{n+1}-k) \gamma_n(p^{n+1}-k).$$

Thus:

$$M_n(\theta, a) = - \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} \frac{(a^{-1})^k}{(a^{-1})^{p^{n+1}}-1} \theta(p^{n+1}-k) \omega^{-1}(p^{n+1}-k) \gamma_n(p^{n+1}-k).$$

The Lemma follows. $\diamond$

Theorem 3.3

$M'(T, \theta, a) \equiv 0 \pmod{\pi_L}$ if and only if $a \equiv -1 \pmod{\pi_L}$ and θ is odd.

Proof By Lemma 3.2, $M(T, \theta, -1) = 0$ if θ is odd. Thus, if $a \equiv -1 \pmod{\pi_L}$ and if θ is odd, we have $M'(T, \theta, a) \equiv 0 \pmod{\pi_L}$.

The proof of this Theorem is based on Sinnott's proof that the Iwasawa μ -invariant vanishes for cyclotomic $\mathbb{Z}_p$ -extensions of abelian number fields ([10]) as exposed in Washington's book([12], paragraph 16.2).

Now, let's suppose that $M'(T, \theta, a) \equiv 0 \pmod{\pi_L}$. We have:

$$\forall n \geq 0, M(T, \theta, a) \equiv \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} \frac{a^k}{a^{p^{n+1}}-1} \theta(k) \omega^{-1}(k) (1+T)^{i(k)} \pmod{\omega_n(T)},$$

where $i(k) = \text{Log}_p(k)/\text{Log}_p(1+p)$. Thus, for all $n \geq 1$, we have:

$$(1+T)M'(T, \theta, a) \equiv \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} i(k) \frac{a^k}{a^{p^{n+1}}-1} \theta(k) \omega^{-1}(k) (1+T)^{i(k)} \pmod{(p^n, \omega_n(T))}.$$

Therefore, for all $n \geq 1$, we get:

$$\sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} i(k) \frac{a^k}{a^{p^{n+1}}-1} \theta(k) \omega^{-1}(k) (1+T)^{i(k)} \equiv 0 \pmod{(\pi_L, \omega_n(T))}.$$

Recall that $i(k) \equiv i(k') \pmod{p^n}$ if and only if $\frac{\leq k > -1}{p} \equiv \frac{\leq k' > -1}{p} \pmod{p^n}$. Therefore changing $i(k)$ to $\frac{\leq k > -1}{p}$ permutes exponents modulo p^n and do not affect divisibility by π_L . Thus:

$$\forall n \geq 1, \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} \frac{\leq k > -1}{p} \frac{a^k}{a^{p^{n+1}} - 1} \theta(k) \omega^{-1}(k) (1+T)^{\frac{\leq k > -1}{p}} \equiv 0 \pmod{(\pi_L, \omega_n(T))}.$$

Let $\alpha \in \mu_{p-1}$. For $n \geq 1$, set:

$$h_n^\alpha(t) \equiv \sum_{k=1, k \equiv \alpha \pmod{p}}^{p^{n+1}-1} \frac{\alpha^{-1}k - 1}{p} \frac{a^k}{a^{p^{n+1}} - 1} \theta(k) \omega^{-1}(k) (1+T)^{\frac{\alpha^{-1}k-1}{p}} \pmod{(p^n, \omega_n(T))}.$$

Note that:

$$h_{n+1}^\alpha(T) \equiv h_n^\alpha(T) \pmod{(p^n, \omega_n(T))}.$$

Now recall that:

$$\Lambda_L \simeq \lim_{\leftarrow} \frac{\Lambda_L}{(p^n, \omega_n(T))}.$$

Therefore, there exists $h_\alpha(T) \in \Lambda_L$ such that:

$$\forall n \geq 1, h_\alpha(T) \equiv h_n^\alpha(T) \pmod{(p^n, \omega_n(T))}.$$

Thus, we have:

$$\sum_{\alpha \in \mu_{p-1}} h_\alpha(T) \equiv 0 \pmod{\pi_L}.$$

And also:

$$\sum_{\alpha \in \mu_{p-1}} (1+T) h_\alpha((1+T)^p - 1) \equiv 0 \pmod{\pi_L}.$$

Now, note that:

$$(1+T) h_n^\alpha((1+T)^p - 1) \equiv f_n^\alpha((1+T)^{\alpha^{-1}} - 1) \pmod{(p^n, \omega_n(T))},$$

where

$$f_n^\alpha(T) \equiv \sum_{k=1, k \equiv \alpha \pmod{p}}^{p^{n+1}-1} \frac{\alpha^{-1}k - 1}{p} \frac{a^k}{a^{p^{n+1}} - 1} \theta(k) \omega^{-1}(k) (1+T)^k \pmod{(p^n, \omega_n(T))}.$$

For $\alpha \in \mu_{p-1}$, let $s_0(\alpha) \in \{1, \dots, p-1\}$ such that $\alpha \equiv s_0(\alpha) \pmod{p}$. We have:

$$f_n^\alpha(T) \equiv \theta(\alpha)\omega^{-1}(\alpha) \sum_{\ell=0}^{p^n-1} \frac{\alpha^{-1}(s_0(\alpha) + p\ell) - 1}{p} \frac{a^{s_0(\alpha)+p\ell}}{a^{p^{n+1}} - 1} (1+T)^{s_0(\alpha)+p\ell} \pmod{(p^n, \omega_n(T))}.$$

But:

$$\sum_{\ell=0}^{p^n-1} \frac{a^{s_0(\alpha)+p\ell}}{a^{p^{n+1}} - 1} (1+T)^{s_0(\alpha)+p\ell} \equiv \frac{a^{s_0(\alpha)}(1+T)^{s_0(\alpha)}}{a^p(1+T)^p - 1} \pmod{(p^n, \omega_n(T))},$$

and

$$\sum_{\ell=0}^{p^n-1} \ell \frac{a^{s_0(\alpha)+p\ell}}{a^{p^{n+1}} - 1} (1+T)^{s_0(\alpha)+p\ell} \equiv -a^p(1+T)^p \frac{a^{s_0(\alpha)}(1+T)^{s_0(\alpha)}}{(a^p(1+T)^p - 1)^2} \pmod{(p^n, \omega_n(T))}.$$

Set:

$$f_\alpha(T) = \theta(\alpha)\omega^{-1}(\alpha) \frac{a^{s_0(\alpha)}(1+T)^{s_0(\alpha)}}{a^p(1+T)^p - 1} \left(\frac{\alpha^{-1}s_0(\alpha) - 1}{p} - \alpha^{-1} \frac{a^p(1+T)^p}{a^p(1+T)^p - 1} \right).$$

Then:

$$\forall n \geq 1, f_n^\alpha(T) \equiv f_\alpha(T) \pmod{(p^n, \omega_n(T))}.$$

We have obtained:

$$\sum_{\alpha \in \mu_{p-1}} f_\alpha((1+T)^{\alpha^{-1}} - 1) \equiv 0 \pmod{\pi_L}.$$

Observe that for all $\alpha \in \mu_{p-1}$, $f_\alpha(T) \in \Lambda_L \cap L(T)$. Thus, by Sinnott's Lemma ([12], Lemma 16.9), for all $\alpha \in \mu_{p-1}$ there exists $c_\alpha \in O_L$ such that:

$$f_\alpha(T) + f_{-\alpha}((1+T)^{-1} - 1) \equiv c_\alpha \pmod{\pi_L}.$$

Observe that:

$$f_\alpha(T) \equiv \theta(\alpha)\omega^{-1}(\alpha) \frac{a^{s_0(\alpha)}}{a^p - 1} \left(\frac{\alpha^{-1}s_0(\alpha) - 1}{p} - \alpha^{-1} \frac{a^p}{a^p - 1} \right) (1+T)^{s_0(\alpha)} \pmod{(\pi_L, T^p)},$$

and

$$f_{-\alpha}((1+T)^{-1} - 1) \equiv \theta(-\alpha)\omega^{-1}(-\alpha) \frac{a^{p-s_0(\alpha)}}{a^p - 1} \left(\frac{-\alpha^{-1}(p-s_0(\alpha))-1}{p} + \alpha^{-1} \frac{a^p}{a^p - 1} \right) (1+T)^{s_0(\alpha)} \pmod{(\pi_L, T^p)}.$$

Thus, for all $\alpha \in \mu_{p-1}$, we must have:

$$a^{s_0(\alpha)} \left(\frac{\alpha^{-1} s_0(\alpha) - 1}{p} - \alpha^{-1} \frac{a^p}{a^p - 1} \right) \equiv \theta(-1) a^{p-s_0(\alpha)} \left(\frac{-\alpha^{-1} (p-s_0(\alpha)) - 1}{p} + \alpha^{-1} \frac{a^p}{a^p - 1} \right) \pmod{\pi_L}.$$

For $\alpha = 1$, we get:

$$a \frac{a^p}{a^p - 1} \equiv -\theta(-1) a^{p-1} \left(\frac{a^p}{a^p - 1} - 1 \right) \pmod{\pi_L}.$$

Thus:

$$a^2 \equiv -\theta(-1) \pmod{\pi_L}.$$

We obtain $a \equiv -1 \pmod{\pi_L}$ and θ is odd or θ is even and $a^2 \equiv -1 \pmod{\pi_L}$. For the second case, if we consider all the equations obtained when α runs through μ_{p-1} , we obtain that for all $b \in \{1, \dots, p-1\}$, b even, we must have:

$$b \equiv \omega(b) + p \pmod{p^2},$$

and for all $b \in \{1, \dots, p-1\}$, b odd, we must have:

$$b \equiv \omega(b) \pmod{p^2},$$

This leads to a contradiction and the Theorem is proved. $\diamond$

We will need the following Lemma:

lemma 3.4 *There exists $\alpha \in \mu_{p-1}$ such that for all prime numbers ℓ , $\ell \equiv \alpha \pmod{p}$ and $\ell \geq p^2$, we have:*

$$\text{Tr}_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}} \left(\frac{\zeta_\ell^{p+1} + \zeta_\ell^{p-1}}{(\zeta_\ell^p - 1)^2} \right) \not\equiv 0 \pmod{p},$$

where ζ_ℓ is a primitive ℓ th root of unity.

Proof For $a \in \mathbb{Z}$, let $[a]_\ell \in \{0, \dots, \ell-1\}$, such that $a \equiv [a]_\ell \pmod{\ell}$. Let $\Phi_\ell(X)$ be the ℓ th cyclotomic polynomial, then:

$$\Phi_\ell(1) = \ell,$$

$$\Phi'_\ell(1) = \frac{\ell(\ell-1)}{2},$$

and

$$\Phi''_\ell(1) = \frac{\ell(\ell-1)(\ell-2)}{3}.$$

Now:

$$\frac{\Phi'_\ell(X)}{\Phi_\ell(X)} = \sum_{\rho \in \mu_\ell \setminus \{1\}} \frac{1}{X - \rho},$$

and

$$\frac{\Phi''_\ell(X)}{\Phi_\ell(X)} - \left(\frac{\Phi'_\ell(X)}{\Phi_\ell(X)}\right)^2 = - \sum_{\rho \in \mu_\ell \setminus \{1\}} \frac{1}{(X - \rho)^2}.$$

Therefore:

$$Tr_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}}\left(\frac{1}{\zeta_\ell - 1}\right) = \frac{1 - \ell}{2},$$

and

$$Tr_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}}\left(\frac{1}{(\zeta_\ell - 1)^2}\right) = \frac{(\ell - 1)^2}{4} - \frac{(\ell - 1)(\ell - 2)}{3}.$$

Furthermore, if $a \in \mathbb{Z}$, $a \not\equiv 0 \pmod{\ell}$, we have:

$$Tr_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}}\left(\frac{\zeta_\ell^a}{\zeta_\ell - 1}\right) = \frac{\ell + 1}{2} - [a]_\ell.$$

Set:

$$S = Tr_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}}\left(\frac{\zeta_\ell^{p+1} + \zeta_\ell^{p-1}}{(\zeta_\ell^p - 1)^2}\right).$$

Let m be the order of p modulo ℓ , set:

$$a = [1 + p^{m-1}]_\ell \text{ and } b = [1 - p^{m-1}]_\ell.$$

We have:

$$S = Tr_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}}\left(\frac{\zeta_\ell^a + \zeta_\ell^b}{(\zeta_\ell - 1)^2}\right).$$

Since $\ell \geq p^2$, we have $b \geq 3$, and thus:

$$a = \ell + 2 - b.$$

Now:

$$S = Tr_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}}\left(\frac{\zeta_\ell^a - 1}{(\zeta_\ell^p - 1)^2}\right) + Tr_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}}\left(\frac{\zeta_\ell^b - 1}{(\zeta_\ell^p - 1)^2}\right) + 2Tr_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}}\left(\frac{1}{(\zeta_\ell^p - 1)^2}\right).$$

We obtain:

$$S = -b^2 + b(\ell + 2) - \frac{\ell^2 + 6\ell + 5}{6}.$$

Therefore, $S \equiv 0 \pmod{p}$ implies that:

$$\frac{\ell^2 + 2}{3} \in (\mathbb{F}_p)^2.$$

Now observe that the function field $\mathbb{F}_p(T, \sqrt{\frac{T^2+2}{3}})$ has genus zero and thus has exactly $p+1$ places of degree one. The Lemma follows. $\diamond$

Theorem 3.5 *Let $\theta \in \hat{\Delta}$, θ even. There exists $\alpha \in \mu_{p-1}$ such that for all prime numbers ℓ , $\ell \equiv \alpha \pmod{p}$ and $\ell \geq p^2$, we have:*

$$\sum_{\rho \in \mu_\ell \setminus \{1\}} M'(T, \theta, \rho) \not\equiv 0 \pmod{p}.$$

Proof

Let ℓ be a prime number, $\ell \neq p$. Suppose that we have:

$$\sum_{\rho \in \mu_\ell \setminus \{1\}} M'(T, \theta, \rho) \equiv 0 \pmod{p}.$$

For $\alpha \in \mu_{p-1}$, set:

$$f_\alpha(T) = \sum_{\rho \in \mu_\ell \setminus \{1\}} \theta(\alpha) \omega^{-1}(\alpha) \frac{\rho^{s_0(\alpha)} (1+T)^{s_0(\alpha)}}{\rho^p (1+T)^p - 1} \left(\frac{\alpha^{-1} s_0(\alpha) - 1}{p} - \alpha^{-1} \frac{\rho^p (1+T)^p}{\rho^p (1+T)^p - 1} \right).$$

By the proof of Theorem 3.3, we get:

$$\sum_{\alpha \in \mu_{p-1}} f_\alpha((1+T)^{\alpha^{-1}} - 1) \equiv 0 \pmod{p}.$$

Therefore by [12], Lemma 16.9, for all $\alpha \in \mu_{p-1}$ there exists $c_\alpha \in \mathbb{Z}_p$ such that:

$$f_\alpha(T) + f_{-\alpha}((1+T)^{-1} - 1) \equiv c_\alpha \pmod{p}.$$

But:

$$f_1(T) \equiv -(1+T) \text{Tr}_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}} \left(\frac{\zeta_\ell^{p+1}}{(\zeta_\ell^p - 1)^2} \right) \pmod{(p, T^p)},$$

and

$$f_{-1}((1+T)^{-1} - 1) \equiv -(1+T) \text{Tr}_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}} \left(\frac{\zeta_\ell^{p-1}}{(\zeta_\ell^p - 1)^2} \right) \pmod{(p, T^p)}.$$

Thus we get:

$$\text{Tr}_{\mathbb{Q}(\zeta_\ell)/\mathbb{Q}} \left(\frac{\zeta_\ell^{p+1} + \zeta_\ell^{p-1}}{(\zeta_\ell^p - 1)^2} \right) \equiv 0 \pmod{p}.$$

It remains to apply Lemma 3.4. $\diamond$

4 p -adic L-functions

Let $\theta \in \widehat{\Delta}$. We set:

- $f(T, \theta) = 0$ if θ is odd,
- if θ is even, $f(T, \theta)$ is the Iwasawa power series associated to the p -adic L-function $L_p(s, \theta)$ (see [12], paragraph 7.2.).

Recall that if $\theta \neq 1$, $f(T, \theta) \in \Lambda$, and if θ is the trivial character then:

$$(1 - \frac{1+p}{1+T})f(T, \theta) \in \Lambda^*.$$

For $\alpha \in \mu_{p-1}$, we set:

$$\rho_n(\alpha) = \frac{\alpha - \zeta_{p^{n+1}}}{\omega(\alpha - 1)} \in U_n,$$

and

$$\rho_\infty(\alpha) = (\rho_n(\alpha))_{n \geq 0} \in U_\infty.$$

We set:

$$\eta_n = \prod_{\alpha \in \mu_{p-1} \setminus \{1\}} \rho_n(\alpha)^{\alpha^{-1}-1} \in U_n,$$

and

$$\eta_\infty = (\eta_n)_{n \geq 0} \in U_\infty.$$

Lemma 4.1 *Let $\theta \in \widehat{\Delta}$, $\theta \neq 1, \omega$. Then:*

$$U_\infty^{e_\theta} = (\eta_\infty^{e_\theta})^\Lambda.$$

Proof By the proof Theorem 2.3:

$$\phi_n^\theta(\rho_n(\alpha)^{e_\theta}) \equiv M_n(\theta, \alpha^{-1})e_{\theta\omega^{-1}}T_n \pmod{p^{n+1}}.$$

Thus:

$$\phi_n^\theta(\eta_n^{e_\theta}) \equiv \left(\sum_{\alpha \in \mu_{p-1} \setminus \{1\}} (\alpha - 1)M_n(\theta, \alpha) \right) e_{\theta\omega^{-1}}T_n \pmod{p^{n+1}}.$$

But:

$$\sum_{\alpha \in \mu_{p-1} \setminus \{1\}} (\alpha - 1) M_0(\theta, \alpha) = (p-1)\theta(-1)\omega^{-1}(-1) \in \mathbb{Z}_p^*.$$

The Lemma follows. $\diamond$

Let $\theta \in \widehat{\Delta}$, $\theta \neq 1, \omega$. We denote by $G(T, \theta) \in \Lambda^*$ the power series that corresponds to $((\sum_{\alpha \in \mu_{p-1} \setminus \{1\}} (\alpha - 1) M_n(\theta, \alpha)))_{n \geq 0}$. Observe that:

$$\sum_{\alpha \in \mu_{p-1} \setminus \{1\}} (\alpha - 1) M_n(\theta, \alpha) = (p-1)\theta(-1)\omega^{-1}(-1)\gamma_n(p-1) \sum_{\ell=1, \ell \not\equiv 0 \pmod{p}}^{p^n} \theta(\ell)\omega^{-1}(\ell)\gamma_n(\ell).$$

Let $\theta \in \widehat{\Delta}$, $\theta \neq 1$, θ even. By $\pi_n^{e_\theta}$ we mean the unique $(p-1)^2$ th root in U_n which is congruent to 1 modulo π_n of:

$$\prod_{a=1}^{p-1} \left(\frac{\zeta_{p^{n+1}}^{\omega(a)} - 1}{\zeta_{p^{n+1}} - 1} \right)^{(p-1)\theta^{-1}(a)}.$$

We set:

$$\pi_\infty^{e_\theta} = (\pi_n^{e_\theta})_{n \geq 0} \in U_\infty.$$

Theorem 4.2

i) Let $\theta \in \widehat{\Delta}$, θ even and non-trivial. Then:

$$(\pi_\infty^{e_\theta})^{-G(\frac{1+T}{1+p}-1, \theta)} = (\eta_\infty^{e_\theta})^{f(\frac{1+p}{1+T}-1, \theta)}.$$

ii) Let $\theta \in \widehat{\Delta}$, $\theta \neq 1, \omega$. Then:

$$\forall \alpha \in \mu_{p-1} \setminus \{1\}, (\rho_\infty(\alpha)^{e_\theta})^{G(\frac{1+T}{1+p}-1, \theta)} = (\eta_\infty^{e_\theta})^{M(\frac{1+T}{1+p}-1, \theta, \alpha^{-1})}.$$

Proof Recall that:

$$\phi_n^\theta(\rho_n(\alpha)^{e_\theta}) \equiv M_n(\theta, \alpha^{-1}) e_{\theta\omega^{-1}} T_n \pmod{p^{n+1}},$$

and

$$\phi_n^\theta(\eta_n^{e_\theta}) \equiv \left(\sum_{\alpha \in \mu_{p-1} \setminus \{1\}} (\alpha - 1) M_n(\theta, \alpha) \right) e_{\theta\omega^{-1}} T_n \pmod{p^{n+1}}.$$

Thus ii) follows from Lemma 4.1, Lemma 2.4, and Proposition 2.2. Note that:

$$\phi_n^\theta(\pi_n^{e_\theta}) \equiv e_{\theta\omega^{-1}} \frac{1}{\zeta_{p^{n+1}} - 1} \pmod{p^{n+1}}.$$

Let $f(X) = \frac{X^{p^{n+1}} - 1}{X - 1}$, then:

$$(X - 1)Xf'(X) + Xf(X) = p^{n+1}X^{p^{n+1}}.$$

Therefore:

$$\frac{1}{\zeta_{p^{n+1}} - 1} = \frac{1}{p^{n+1}} \sum_{k=1}^{p^{n+1}-1} k \zeta_{p^{n+1}}^k.$$

Let $\theta \in \widehat{\Delta}$, for $\theta \neq 1$, set:

$$v_n(\theta) = -\frac{1}{p^{n+1}} \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} k \theta(k) \omega^{-1}(k) \gamma_n(k),$$

and if $\theta = 1$, set:

$$v_n(\theta) = -(1 - (1 + p)\gamma_n(1 + p)) \frac{1}{p^{n+1}} \sum_{k=1, k \not\equiv 0 \pmod{p}}^{p^{n+1}-1} k \omega^{-1}(k) \gamma_n(k).$$

Then by [12], paragraph 7.2, $(v_n(\theta))_{n \geq 0}$ corresponds to $f(\frac{1}{1+T} - 1, \theta)$ if $\theta \neq 1$, and to $(1 - (1 + p)(1 + T))f(\frac{1}{1+T} - 1, \theta)$ if θ is trivial. Observe that if θ is even and non-trivial:

$$e_{\theta\omega^{-1}} \frac{1}{\zeta_{p^{n+1}} - 1} = -v_n(\theta) e_{\theta\omega^{-1}} T_n,$$

and if $\theta = 1$:

$$(1 - (1 + p)\gamma_n(1 + p)) e_{\theta\omega^{-1}} \frac{1}{\zeta_{p^{n+1}} - 1} = -v_n(\theta) e_{\theta\omega^{-1}} T_n.$$

The Theorem follows. $\diamond$

Theorem 4.3 *Let d be an integer, $d \geq 2$, $d \not\equiv 0 \pmod{p}$. Let $\theta \in \widehat{\Delta}$. Then:*

$$\sum_{\rho \in \mu_d, o(\rho)=d} M(T, \theta, \rho) = -f\left(\frac{1}{1+T} - 1, \theta\right) \left(\sum_{\ell \text{ divides } d} \ell \mu\left(\frac{d}{\ell}\right) \theta(\ell) \omega^{-1}(\ell) (1+T)^{\frac{\text{Log}_p(\ell)}{\text{Log}_p(1+p)}} \right),$$

wher $\mu(\cdot)$ is the Möbius function.

Proof If θ is odd, the result is clear by Lemma 3.2. Thus we assume that θ is even. Let $\Phi_d(X)$ be the d th cyclotomic polynomial, i.e.

$$\Phi_d(X) = \prod_{\rho \in \mu_d, o(\rho)=d} (X - \rho) \in \mathbb{Z}[X].$$

Then:

$$\Phi_d(X) = \prod_{\ell \text{ divides } d} (X^\ell - 1)^{\mu(d/\ell)}.$$

If we take logarithmic derivatives, we get:

$$\frac{\Phi'_d(X)}{\Phi_d(X)} = \sum_{\ell \text{ divides } d} \ell \mu\left(\frac{d}{\ell}\right) \frac{X^{\ell-1}}{X^\ell - 1}.$$

It follows that:

$$\sum_{\rho \in \mu_d, o(\rho)=d} \frac{\zeta_{p^{n+1}}}{\zeta_{p^{n+1}} - \rho} = \sum_{\ell \text{ divides } d} \ell \mu\left(\frac{d}{\ell}\right) \frac{\zeta_{p^{n+1}}^\ell}{\zeta_{p^{n+1}}^\ell - 1}.$$

Thus:

$$\sum_{\rho \in \mu_d, o(\rho)=d} \frac{1}{\rho \zeta_{p^{n+1}} - 1} = \left(\sum_{\ell \text{ divides } d} \ell \mu\left(\frac{d}{\ell}\right) \sigma_n(\ell) \right) \frac{1}{\zeta_{p^{n+1}} - 1}.$$

Now:

$$\frac{\rho^{p^{n+1}} - 1}{\rho \zeta_{p^{n+1}} - 1} - 1 = \sum_{k=1}^{p^{n+1}-1} \rho^k \zeta_{p^{n+1}}^k.$$

Thus:

$$\frac{1}{\rho \zeta_{p^{n+1}} - 1} - \frac{1}{\rho^{p^{n+1}} - 1} = \sum_{k=1}^{p^{n+1}-1} \frac{\rho^k}{\rho^{p^{n+1}} - 1} \zeta_{p^{n+1}}^k.$$

We are working in the extension $\mathbb{Q}_p(\mu_d, \zeta_{p^{n+1}})/\mathbb{Q}_p(\mu_d)$ and we identify $\text{Gal}(\mathbb{Q}_p(\mu_d, \zeta_{p^{n+1}})/\mathbb{Q}_p(\mu_d))$ with $\Gamma_n \times \Delta$. Therefore:

$$e_{\theta\omega^{-1}} \frac{1}{\rho \zeta_{p^{n+1}} - 1} = M_n(\theta, \rho) e_{\theta\omega^{-1}} \zeta_{p^{n+1}} + e_{\theta\omega^{-1}} \left(\sum_{k=1}^{p^{n+1}-1} \frac{(\rho^p)^k}{(\rho^p)^{p^n} - 1} \zeta_{p^n}^k \right).$$

Now observe that the map $\mu_d \rightarrow \mu_d, \rho \mapsto \rho^p$, is an isomorphism. Thus:

$$e_{\theta\omega^{-1}} \left(\sum_{\rho \in \mu_d, o(\rho)=d} \frac{1}{\rho \zeta_{p^{n+1}} - 1} \right) = \left(\sum_{\rho \in \mu_d, o(\rho)=d} M_n(\theta, \rho) \right) e_{\theta\omega^{-1}} T_n.$$

But T_n generates a normal basis for the field extension $K_n/\mathbb{Q}_p$, thus:

$$\sum_{\rho \in \mu_d, o(\rho)=d} M_n(\theta, \rho) = - \left(\sum_{\ell \text{ divides } d} \ell \mu\left(\frac{d}{\ell}\right) \theta(\ell) \omega^{-1}(\ell) \gamma_n(\ell) \right) v_n(\theta),$$

where $v_n(\theta)$ is as in the proof of Theorem 4.2. The Theorem follows. $\diamond$

Corollary 4.4 *Let $\theta \in \widehat{\Delta}$, θ even and non-trivial. Then:*

$$f'(T, \theta) \not\equiv 0 \pmod{p}.$$

Proof Let $\alpha \in \mu_{p-1}$ as in Theorem 3.5. Let ℓ be a prime number such that $\ell \geq p^2$ and $\ell \equiv \alpha \pmod{p^2}$ (note that there exist infinitely many such primes). We know that:

$$\sum_{\rho \in \mu_\ell \setminus \{1\}} M'(T, \theta, \rho) \not\equiv 0 \pmod{p}.$$

But by Theorem 4.3:

$$\sum_{\rho \in \mu_\ell \setminus \{1\}} M(T, \theta, \rho) = -f\left(\frac{1}{1+T} - 1, \theta\right) (\ell \theta(\ell) \omega^{-1}(\ell) (1+T)^{\frac{\text{Log}_p(\ell)}{\text{Log}_p(1+p)}} - 1).$$

But since $\ell^{p-1} \equiv 1 \pmod{p^2}$, we have:

$$\frac{\text{Log}_p(\ell)}{\text{Log}_p(1+p)} \equiv 0 \pmod{p}.$$

Thus, if we take derivatives and reduce modulo p , we get:

$$\sum_{\rho \in \mu_\ell \setminus \{1\}} M'(T, \theta, \rho) \equiv \frac{1}{(1+T)^2} f'\left(\frac{1}{1+T} - 1, \theta\right) (\ell \theta(\ell) \omega^{-1}(\ell) (1+T)^{\frac{\text{Log}_p(\ell)}{\text{Log}_p(1+p)}} - 1) \pmod{p}.$$

The Corollary follows. $\diamond$

The case of the trivial character is treated in the last section.

5 Other results

Let θ be an even Dirichlet character of conductor d or pd , where $d \geq 1$, $d \not\equiv 0 \pmod{p}$. For all $n \geq 0$, set: $q_n = p^{n+1}d$. Let $g(T, \theta)$ be the power series introduced in [12], paragraph 7.2, i.e.

$$g(T, \theta) = \frac{T - q_0}{1 + T} f(T, \theta),$$

where $f(T, \theta)$ is the power series associated to the p -adic L-function $L_p(s, \theta)$ (see [12], Theorem 7.10). Set $L = \mathbb{Q}_p(\theta)$, and let π_L be a prime of L . Then, the Ferrero-Washington Theorem states (see [12], paragraph 16.2):

$$g(T, \theta) \not\equiv 0 \pmod{\pi_L}.$$

We have:

Theorem 5.1

$$g'(T, \theta) \not\equiv 0 \pmod{\pi_L}.$$

Proof For $y \in \mathbb{Q}$, set:

$$B(y) = (1 + q_0)\{y\} - \{(1 + q_0)y\} - \frac{q_0}{2} \in \mathbb{Z}_p,$$

where $\{y\}$ is the fractional part of y . recall that ([12], proof of Theorem 7.10):

$$\forall n \geq 0, g(T, \theta) \equiv \sum_{a=1, (a, q_0)=1}^{q_n} B\left(\frac{a}{q_n}\right) \theta \omega^{-1}(a) (1 + T)^{-i(a)-1} \pmod{\omega_n(T)},$$

where $i(a) = \frac{\text{Log}_p(a)}{\text{Log}_p(1+q_0)}$. Let's suppose that $g'(T, \theta) \equiv 0 \pmod{p}$. We have, for all $n \geq 1$:

$$(1+T)^2 g'(T, \theta) \equiv - \sum_{a=1, (a, q_0)=1}^{q_n} (i(a)+1) B\left(\frac{a}{q_n}\right) \theta \omega^{-1}(a) (1+T)^{-i(a)} \pmod{(p^n, \omega_n(T))}.$$

Therefore, we get:

$$\forall n \geq 1, \sum_{a=1, (a, q_0)=1}^{q_n} (i(a)+1) B\left(\frac{a}{q_n}\right) \theta \omega^{-1}(a) (1+T)^{i(a)} \equiv 0 \pmod{(\pi_L, \omega_n(T))}.$$

Now, changing $i(a)$ to $(1 + q_0)^{\frac{\langle a \rangle - 1}{p}}$ permutes exponents modulo p^n and does not affect divisibility by π_L . Thus, for all $n \geq 1$:

$$\sum_{a=1, (a, q_0)=1}^{q_n} ((1+q_0)^{\frac{\langle a \rangle - 1}{p}} + 1) B\left(\frac{a}{q_n}\right) \theta \omega^{-1}(a) (1+T)^{(1+q_0)^{\frac{\langle a \rangle - 1}{p}}} \equiv 0 \pmod{(\pi_L, \omega_n(T))}.$$

For $n \geq 1$ and for $\alpha \in \mu_{p-1}$, set:

$$H_\alpha^n(T) \equiv \sum_{a \equiv \alpha \pmod{p}} ((1+q_0)^{\frac{\alpha^{-1}a-1}{p}} + 1) B\left(\frac{a}{q_n}\right) \theta \omega^{-1}(a) (1+T)^{(1+q_0)^{\frac{\alpha^{-1}a-1}{p}}} \pmod{(p^n, \omega_n(T))}.$$

Note that:

$$H_\alpha^{n+1}(T) \equiv H_\alpha^n(T) \pmod{(p^n, \omega_n(T))}.$$

Thus, there exists $H_\alpha(T) \in \Lambda_L$ such that:

$$H_\alpha(T) \equiv H_\alpha^n(T) \pmod{(p^n, \omega_n(T))}.$$

We get:

$$\sum_{\alpha \in \mu_{p-1}} H_\alpha(T) \equiv 0 \pmod{\pi_L}.$$

Therefore:

$$\sum_{\alpha \in \mu_{p-1}} (1+T)^{1+q_0} H_\alpha((1+T)^p - 1) \equiv 0 \pmod{\pi_L}.$$

Let $f_\alpha(T) \in \Lambda_L$ as in [12], Lemma 16.8. Set:

$$F_\alpha(T) = \frac{\alpha^{-1}}{p} (1+T) f'_\alpha(T) + \left(1 - \frac{1+q_0}{p}\right) f_\alpha(T).$$

It is not difficult to see that $F_\alpha(T) \in \Lambda_L \cap L(T)$. By [12], Lemma 16.8, we have:

$$F_\alpha(T) \equiv \sum_{a \equiv \alpha \pmod{p}} ((1+q_0)^{\frac{\alpha^{-1}a-1}{p}} + 1) B\left(\frac{a}{q_n}\right) \theta \omega^{-1}(a) (1+T)^{(1+q_0)a} \pmod{(p^n, \omega_n(T))}.$$

Therefore, we get:

$$\sum_{\alpha \in \mu_{p-1}} F_\alpha((1+T)^{\alpha^{-1}} - 1) \equiv 0 \pmod{\pi_L}.$$

Now apply [12], Lemma 16.9, and we get that for all $\alpha \in \mu_{p-1}$, there exists $b_\alpha \in O_L$ such that:

$$F_\alpha(T) + F_{-\alpha}((1+T)^{-1} - 1) \equiv b_\alpha \pmod{\pi_L}.$$

But observe that:

$$F_\alpha(T) = F_{-\alpha}((1+T)^{-1} - 1).$$

Thus, for all $\alpha \in \mu_{p-1}$, there exists $c_\alpha \in O_L$ such that:

$$\frac{\alpha^{-1}}{p}(1+T)f'_\alpha(T) + (1 - \frac{1+q_0}{p})f_\alpha(T) \equiv c_\alpha \pmod{\pi_L}.$$

Now we take $\alpha = 1$, and we set:

$$G_1(T) = (1+q_0) \sum_{0 < a < q_0, a \equiv 1 \pmod{p}} \theta \omega^{-1}(a) (1+T)^{a(1+q_0)} - \sum_{0 < a < q_0^2 + q_0, a \equiv 1 \pmod{p}} \theta \omega^{-1}(a) (1+T)^a.$$

We get:

$$(1 - \frac{1+q_0}{p})((1+T)^{q_0(1+q_0)} - 1)G_1(T) + \frac{1}{p}(1+T)((1+T)^{q_0(1+q_0)} - 1)G'_1(T) - q_0(1+q_0)(1+T)^{q_0(1+q_0)-1}G_1(T) \equiv c_1((1+T)^{q_0(1+q_0)} - 1)^2 \pmod{\pi_L}.$$

Note that:

$$G_1(T) \equiv -(1+T) \pmod{(1+T)^{1+p}},$$

ans

$$G'_1(T) \equiv -1 \pmod{(1+T)^p}.$$

Thus we must have:

$$c_1 \equiv 0 \pmod{\pi_L}.$$

The coefficient of $1+T$ in the left-side is: $1-d$ and the coefficient of $(1+T)^{q_0(1+q_0)+1}$ in the left side is $2d-1+\frac{q_0^2}{p}$. The Theorem follows. $\diamond$

References

- [1] A. Adler and L. C. Washington, p -adic L -functions and higher-dimensional magic cubes, J. Number Theory **52** (1995), 179-197.
- [2] B. Anglès, Units and norm residue symbol, Acta Arith. **98** (2001), 33-51.
- [3] R. Coleman, Division values in local fields, Invent. Math. **53** (1979), 91-116.
- [4] B. Ferrero and L. C. Washington, The Iwasawa invariant μ_p vanishes for abelian number fields, Annals of Math. **109** (1979), 377-395.
- [5] S. Lang, Cyclotomic Fields I & II, Springer-Verlag (1990).

- [6] H. W. Leopoldt, Über die Hauptordnung der ganzen Elemente eines abelschen Zahlkörpers, J. Reine angew. Math. **201** (1959), 119-149.
- [7] G. Lettl, The ring of integers of an abelian number field, J. Reine angew. Math. **404** (1990), 162-170.
- [8] T. Metsänkylä, Iwasawa invariants and Kummer congruences, J. Number Theory **10** (1978), 510-522.
- [9] P. Ribenboim, 13 Lectures on Fermat's Last Theorem, Springer-Verlag (1979).
- [10] W. Sinnott, On the μ -invariant of the Γ -transform of a rational function, Invent. Math. **75** (1984), 273-282.
- [11] G. Terjanian, Sur la loi de réciprocité des puissances ℓ -ièmes, Acta Arith. **54** (1989), 87-125.
- [12] L. C. Washington, Introduction to Cyclotomic Fields, Second Edition, Springer-Verlag (1997).